

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show \\$ Numbers](#)[Edit \\$ Numbers](#)[Preferences](#)

09/17/916
 Search results
 for Page # 16

Search Results -

Terms	Documents
APC\$ near10 RNA near10 tumor\$	3

Database:

US Patents Full-Text Database
 US Pre-Grant Publication Full-Text Database
 JPO Abstracts Database
 EPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Refine Search:

APC\$ near10 RNA near10 tumor\$

[Clear](#)**Search History****Today's Date: 11/4/2001**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	APC\$ near10 RNA near10 tumor\$	3	<u>L8</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	APC\$ near5 RNA near5 tumor\$	3	<u>L7</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	14 and APC near5 RNA near5 tumor\$	1	<u>L6</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	14 and APC	1	<u>L5</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	6306640 [pn]	1	<u>L4</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	cytotoxic near T near lymphocyte\$ near5 immunotherap\$ near10 tumor\$	8	<u>L3</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	cytotoxic near T near lymphocyte\$ near5 immunotherap\$	30	<u>L2</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	CTL\$ near5 clon\$ near5 isolat\$ near5 antigen\$	10	<u>L1</u>

WEST

Generate Collection

L1: Entry 6 of 10

File: USPT

Oct 5, 1999

DOCUMENT-IDENTIFIER: US 5962318 A

TITLE: Cytotoxic T lymphocyte-mediated immunotherapy

BSPR:

In such a situation, one may wish to isolate and expand the CTL clone(s) that specifically recognizes the subdominant antigen of choice, e.g., LMP-2a, or one may wish to develop methods for culturing CTLs in vitro that only recognize LMP-2a, i.e., they are monospecific. In order to maximize the T cell response to a weak immunogen, the target antigen would need to be expressed on the surface of a powerful antigen presenting cell. Thus, there is a need in the art to develop a system that provides for high level presentation of the weak antigen on an effective antigen presenting cell. There is a further need to generate antigen specific CTLs to induce both primary and memory T cell responses to a subdominant antigen. There is still a further specific need to develop a transduction system that provides for high level expression of a foreign gene in antigen presenting cells, and application of such a system to generate CTLs for the treatment of malignancies expressing specific viral or tumor antigens.

WEST

Generate Collection

L2: Entry 19 of 30

File: USPT

Oct 27, 1998

DOCUMENT-IDENTIFIER: US 5827642 A

TITLE: Rapid expansion method ("REM") for in vitro propagation of T lymphocytes

ABPL:

The present invention provides a rapid expansion method (termed "REM"), for quickly generating large numbers of T lymphocytes, including cytolytic and helper T lymphocytes. REM involves culturing the T cells in association with a disproportionately large concentration of nondividing feeder cells, preferably .gamma.-irradiated peripheral blood mononuclear cells ("PBMC") present at an excess of at least 40-fold (relative to the number of target T cells), more preferably at an excess of at least about 200-fold. Cultures grown under REM exhibit dramatically enhanced expansion rates that can be even further elevated by the use of appropriate concentrations of an additional feeder cell, an anti-CD3 monoclonal antibody and IL-2, as described herein. Clonal expansions in the range of 500-fold to 3000-fold can be achieved within a single stimulation cycle of about 10-13 days, which is more than 100-fold more efficient than currently employed methods of culturing human T cell clones. Genetic transduction efficiencies were also enhanced using REM-expanded T lymphocytes. Several examples involving human bone marrow transplant recipients illustrate the effective use of REM-expanded antigen-specific cytotoxic T lymphocytes for adoptive immunotherapy in humans.

WEST☐ Generate Collection

L3: Entry 1 of 8

File: USPT

Aug 21, 2001

DOCUMENT-IDENTIFIER: US 6277368 B1

TITLE: Cancer immunotherapy using autologous tumor cells
combined with cells expressing a membrane cytokine

ORPL:

Kruse et al., "Analysis of interleukin 2 and various effector
cell populations in adoptive immunotherapy of 9L rat
gliosarcoma: Allogeneic cytotoxic T lymphocytes prevent tumor
take" Proc. Natl. Acad. Sci. USA (1990) 87:9577-9581.